

Macroscopic regularity for the Boltzmann equation

Feimin Huang^a, Yong Wang^{a*},

^a*Institute of Applied Mathematics, AMSS, CAS, Beijing 100190, China*

Abstract

The regularity of solutions to the Boltzmann equation is a fundamental problem in the kinetic theory. In this paper, the case with angular cut-off is investigated. It is shown that the macroscopic parts of solutions to the Boltzmann equation, i.e. the density, momentum and total energy are continuous functions of (x, t) in the region $\mathbb{R}^3 \times (0, +\infty)$. More precisely, these macroscopic quantities immediately become continuous in any positive time even though they are initially discontinuous and the discontinuities of solutions propagate only in the microscopic level. It should be noted that such kind of phenomenon can not happen for the compressible Navier-Stokes equations in which the initial discontinuities of the density never vanish in any finite time, see [20]. This hints that the Boltzmann equation has better regularity effect in the macroscopic level than compressible Navier-Stokes equations.

Keywords: Boltzmann equation, Macroscopic regularity, Compressible Navier-Stokes equations

AMS: 35Q35, 35B65, 76N10

1 Introduction

In this paper, we consider the Boltzmann equation

$$F_t + v \cdot \nabla_x F = Q(F, F), \quad (1.1)$$

where $F(t, x, v)$ is a distribution function for the gas particles at time $t > 0$, position $x \in \mathbb{R}^3$ and particle velocity $v \in \mathbb{R}^3$. The collision operator $Q(F, F)$ takes the following bilinear form

$$\begin{aligned} Q(F_1, F_2) &= \int_{\mathbb{R}^3} \int_{\mathbb{S}^2} q(\theta, v - u) F_1(u') F_2(v') du d\omega - \int_{\mathbb{R}^3} \int_{\mathbb{S}^2} q(\theta, v - u) F_1(u) F_2(v) du d\omega \\ &\triangleq Q_+(F_1, F_2) - Q_-(F_1, F_2), \end{aligned} \quad (1.2)$$

where the relation between the post-collision velocity (v', u') of the two particles with pre-collision velocity pair (v, u) is given by

$$u' = u + [(v - u) \cdot \omega] \omega, \quad v' = v - [(v - u) \cdot \omega] \omega,$$

satisfying

$$u' + v' = u + v, \quad \text{and} \quad |u'|^2 + |v'|^2 = |u|^2 + |v|^2.$$

*Corresponding author.

Email addresses: fhuang@amt.ac.cn(Feimin Huang), yongwang@amss.ac.cn(Yong Wang)

Throughout this paper, we consider the hard and soft potentials with angular cut-off, i.e.,

$$q(\theta, v - u) = |v - u|^\gamma b(\theta), \text{ and } \int_0^{\frac{\pi}{2}} b(\theta) d\theta < +\infty, \text{ with } -3 < \gamma \leq 1. \quad (1.3)$$

We impose the Boltzmann equation (1.1) with initial condition

$$F(t, x, v)|_{t=0} = F_0(x, v) \geq 0. \quad (1.4)$$

There is an enormous literature on the study of well-posedness of the Cauchy problem for the Boltzmann equation. The existence of global renormalized solutions was first proved by R. Diperna and P.-L. Lions [12] for large initial data. However, little is known on the uniqueness of renormalized solution. For small initial data, various results on the existence and uniqueness of solutions were obtained, see [5, 11, 17, 18, 19, 21, 22, 23, 25, 28, 29] and references therein. We also refer to the monographs [4, 10, 15] and survey paper [30].

On the other hand, the regularity of solutions is a key problem of the Boltzmann equation. For the case with angular cut-off, it is believed that the initial singularities propagate in time since the Boltzmann equation is hyperbolic. This property was proved by Boudin-Desvillettes [7] with propagation of Sobolev $H^{\frac{1}{25}}$ singularity in the case near vacuum, later by Duan-Li-Yang [14] in the case near global Maxwellian. We also refer to [3, 6, 24, 27] for other interesting works. For the case without angular cut-off, the solution is believed more regular since the non-cutoff collision operator behaves like a fractional Laplacian. For interesting results on this topic, we refer to [1, 2, 8, 9, 13, 26] and the references therein.

It is well known that the Boltzmann equation is closely related to the compressible Navier-Stokes equations through Chapman-Enskog expansion. It is conjectured that the Boltzmann equation has regularity in the macroscopic components. In fact, the famous average lemma [16] shows that $\int_{\mathbb{R}^3} f(t, x, v) \varphi(v) dv \in H^{\frac{1}{2}}(t, x)$ for any smooth compact support function $\varphi(v)$. This indicates that the macroscopic components like density, momentum and total energy probably have $H^{\frac{1}{2}}(t, x)$ regularity. However more regularity is not known from the average lemma. The purpose of this paper is to investigate better regularity for the macroscopic quantities of solutions to Boltzmann equation (1.1) with angular cut-off. We find that the Boltzmann equation has better macroscopic regularity beyond the compressible Navier-Stokes equations. Precisely speaking, we shall prove that the macroscopic parts of solutions to the Boltzmann equation, i.e. the density, momentum and total energy are continuous functions of (x, t) in the region $\mathbb{R}^3 \times (0, +\infty)$. In other words, these macroscopic quantities immediately become continuous in any positive time even though they are initially discontinuous and the discontinuities of solutions propagate only in the microscopic level. It should be noted that such kind of phenomenon can not happen for the compressible Navier-Stokes equations in which the initial discontinuities of the density never vanish in any finite time, see [20]. This means that the Boltzmann equation has better regularity effect in the macroscopic level beyond the compressible Navier-Stokes equations.

Define a weight function

$$w(v) = (1 + |v|^2)^{\frac{\beta}{2}}, \quad (1.5)$$

and let the global Maxwellian μ be

$$\mu(v) = \frac{1}{(2\pi)^{\frac{3}{2}}} \exp\left(-\frac{|v|^2}{2}\right). \quad (1.6)$$

Let F_0 be the initial data satisfying, for any fixed $(x, t) \in \mathbb{R}^3 \times (0, +\infty)$,

$$\|wF_0\|_{L_{x,v}^\infty} < +\infty \text{ and } \int_{\mathbb{R}_{loc}^3} |F_0(x + \delta - v(t + \sigma), v) - F_0(x - vt, v)| dv \rightarrow 0, \text{ as } \sigma, \delta \rightarrow 0. \quad (1.7)$$

We note that the initial condition (1.7) is general. Indeed, there is a large class of initial data satisfying (1.7). For the case near global Maxwellian, we can choose, for example, that

$$F_0(x, v) = \frac{\rho_0(x)}{(2\pi\theta_0(x))^{\frac{3}{2}}} (1 + P(v)) \exp\left(-\frac{|v - u_0(x)|^2}{2\theta_0(x)}\right), \quad (1.8)$$

where $|P(v)| \leq C(1 + |v|)^m$ holds for some positive constant $m > 0$, which is allowed to be discontinuous in $v \in \mathbb{R}^3$, ρ_0, θ_0, u_0 may also be discontinuous in $x \in \mathbb{R}^3$ and satisfy

$$0 \leq \rho_0(x) \leq \hat{C} < +\infty, \quad \hat{C}^{-1} \leq \theta_0(x) \leq \hat{C} < +\infty, \quad |u_0(x)| \leq \hat{C} < +\infty, \quad (1.9)$$

with

$$\|(\rho_0 - 1, \theta_0 - 1, u_0)\|_{L^p} < +\infty, \text{ for some } p \in [1, \infty). \quad (1.10)$$

For the case near vacuum, one can choose, for example, that

$$F_0(x, v) = \frac{1}{(2\pi)^{\frac{3}{2}}} \rho_0(x) (1 + P(v)) \exp\left(-\frac{|v - u_0(x)|^2}{2}\right), \quad (1.11)$$

where ρ_0 and u_0 satisfy

$$\begin{cases} 0 \leq \rho_0(x) \leq \hat{C} < +\infty, & |u_0(x)| \leq \hat{C} < +\infty, \\ \|(\rho_0, u_0)\|_{L^p} < +\infty, & \text{for some } p \in [1, \infty). \end{cases} \quad (1.12)$$

One can check that both (1.8) and (1.11) satisfy the condition (1.7), see appendix below.

The solutions considered in this paper are in the following space $\mathbf{X}$:

Near Maxwellian μ , $F(t, x, v) \geq 0$ for a.e. $(t, x, v) \in (0, +\infty) \times \mathbb{R}^3 \times \mathbb{R}^3$ and

$$w(\cdot)[F(\cdot, \cdot, \cdot) - \mu(\cdot)] \in L^{\frac{b}{b-1}}(0, T; L_{x,v}^2) \text{ and } w(\cdot)F(\cdot, \cdot, \cdot) \in L^b(0, T; L_{x,v}^\infty) \text{ for } b > 2; \quad (1.13)$$

Near vacuum, $F(t, x, v) \geq 0$ for a.e. $(t, x, v) \in (0, +\infty) \times \mathbb{R}^3 \times \mathbb{R}^3$ and

$$w(\cdot)F(\cdot, \cdot, \cdot) \in L^{\frac{b}{b-1}}(0, T; L_{x,v}^2) \text{ and } w(\cdot)F(\cdot, \cdot, \cdot) \in L^b(0, T; L_{x,v}^\infty) \text{ for } b > 2. \quad (1.14)$$

Then our main results are as follows:

Theorem 1.1 *Let (1.3) hold, $\beta > 8 + 2\max\{0, \gamma\}$, the initial data F_0 satisfy (1.7), and the solutions $F(t, x, v)$ of Boltzmann equation belong to the space $\mathbf{X}$, then the macroscopic components of solutions to the Boltzmann equation (1.1) are continuous functions of (x, t) in the region $\mathbb{R}^3 \times (0, +\infty)$.*

Moreover, we have

Theorem 1.2 *Under the conditions of Theorem 1.1, and if the initial data F_0 further satisfies*

$$\sup_{(x,t) \in \mathbb{R}^3 \times [t_1, T]} \int_{\mathbb{R}_{loc}^3} |F_0(x + \delta - v(t + \sigma), v) - F_0(x - vt, v)| dv \rightarrow 0, \text{ as } \sigma, \delta \rightarrow 0. \quad (1.15)$$

where t_1 and T are any given times with $0 < t_1 < T < +\infty$, then the macroscopic components of solutions to the Boltzmann equation (1.1) are uniformly continuous functions of (x, t) in $\mathbb{R}^3 \times [t_1, T]$.

A few remarks are in order.

Remark 1.3 *It is noted that the condition (1.15) is slightly stronger than (1.7). Indeed, both examples in (1.8) and (1.11) satisfy (1.15) for any $0 < t_1 < T < +\infty$, see the appendix below.*

Remark 1.4 *It is noted that even though the initial macroscopic parts in Theorems 1.1 and 1.2 are allowed to be discontinuous in $x \in \mathbb{R}^3$ (the possible discontinuous set has zero measure), the macroscopic components of solutions to the Boltzmann equation immediately become continuous in (x, t) for any positive time $t > 0$. Such kind of phenomenon can not happen for the compressible Navier-Stokes equations in which the initial discontinuities of the density never vanish in any finite time, see [20]. This hints that the Boltzmann equation has better regularity effect in the macroscopic level beyond the compressible Navier-Stokes equations, and the discontinuities of solutions to the Boltzmann equation propagate only in the microscopic level.*

Remark 1.5 *Theorems 1.1 and 1.2 are still valid if the solution space $L^{\frac{b}{b-1}}(0, T; L^2_{x,v})$ is relaxed to $L^{\frac{b}{b-1}}(0, T; L^a_{x,v})$ with $a > 1$.*

Remark 1.6 *If $\beta > 8 + 2 \max\{0, \gamma\} + k$, $k \in \mathbb{R}_+$, let $B(v)$ be any function satisfying $|B(v)| \leq C(1 + |v|)^{2+k}$, then, by the same arguments as in the proof of Theorems 1.1 and 1.2, one can still prove that*

$$\int_{\mathbb{R}^3} B(v) F(t, x, v) dv,$$

has the same results as in Theorems 1.1 and 1.2.

Remark 1.7 *It is noted that the solutions constructed in [29, 18] near Maxwellian and [5, 19, 21, 23, 25] near vacuum belong to $\mathbf{X}$. If their initial data further satisfy (1.7) (or (1.15)), then Theorem 1.1 (or Theorem 1.2) holds. That is, the macroscopic quantities of their solutions are continuous in (x, t) . Thus, Theorems 1.1 and 1.2 are self-contained in this sense.*

The proof of Theorem 1.1 and Theorem 1.2 is based on the following observation. Since the Boltzmann equation is essentially transport, the solution propagates along the velocity v direction. Thus the integral with respect to v can be translated to the integral with respect to x by the change of variables. The continuity of macroscopic components is then derived from the continuity of translations on $L^p_{x,t}$ space. The formula of collision operator plays important role in the whole proof.

The rest of this paper is arranged as follows. Section 2 is devoted to the proof of Theorem 1.1. The proof of Theorem 1.2 is given in section 3.

Notations. Throughout this paper, C denotes a generic positive constant which may depend on β, γ and vary from line to line. $C(t)$ denotes the generic positive continuous function depending on time $t > 0$ and β, γ which also may vary from line to line. $\|\cdot\|_{L^2}$ denotes the standard $L^2(\mathbb{R}^3_x \times \mathbb{R}^3_v)$ -norm, and $\|\cdot\|_{L^\infty}$ denotes the $L^\infty(\mathbb{R}^3_x \times \mathbb{R}^3_v)$ -norm.

2 Proof of the Main Theorems

Let $F(t, x, v)$ be the solution of Boltzmann equation (1.1), we write down the Duhamel form of the solution to Boltzmann equation (1.1). For $(t, x, v) \in (0, +\infty) \times \mathbb{R}^3 \times \mathbb{R}^3$, one has that

$$\begin{aligned} F(t, x, v) &= F_0(x - vt, v) e^{-\int_0^t g(\tau; t, x, v) d\tau} + \int_0^t e^{-\int_s^t g(\tau; t, x, v) d\tau} Q_+(F, F)(s, x - v(t - s), v) ds \\ &\triangleq I_1(t, x, v) + I_2(t, x, v), \end{aligned} \tag{2.1}$$

where

$$g(\tau; t, x, v) = \int_{\mathbb{R}^3} \int_{\mathbb{S}^2} q(\theta, v - u) F(\tau, x - v(t - \tau), u) d\omega du \geq 0. \quad (2.2)$$

Proof of Theorem 1.1. For any fixed $(t, x) \in (0, +\infty) \times \mathbb{R}^3$, let $\sigma \in \mathbb{R}$, $\delta \in \mathbb{R}^3$ with $|\sigma| \leq \min\{1, \frac{t}{2}\}$ and $|\delta| \leq 1$. The continuity of density $\rho(x, t)$ in (x, t) is equivalent to show that

$$\left| \int_{\mathbb{R}^3} F(t + \sigma, x + \delta, v) - F(t, x, v) dv \right| \rightarrow 0 \text{ as } \sigma, \delta \rightarrow 0. \quad (2.3)$$

That is, for any $\varepsilon > 0$, we need to prove that there exists $\chi > 0$, which may depend on (x, t) and ε , such that if $|\sigma| \leq \chi$ and $|\delta| \leq \chi$,

$$\left| \int_{\mathbb{R}^3} F(t + \sigma, x + \delta, v) - F(t, x, v) dv \right| \leq \varepsilon. \quad (2.4)$$

We assume $\sigma \geq 0$ without loss of generality and denote $t_\sigma \doteq t + \sigma$ for notation simplicity. We divide the proof into two parts. The proof is based on the initial condition (1.7) and the continuity of translations on L^p .

Part 1: Estimation on $\int_{\mathbb{R}^3} I_1(t, x, v) dv$.

Firstly, we note that

$$\begin{aligned} J_1 &\triangleq \left| \int_{\mathbb{R}^3} (I_1(t_\sigma, x + \delta, v) - I_1(t, x, v)) dv \right| \\ &\leq \int_{\mathbb{R}^3} \left| F_0(x + \delta - vt_\sigma, v) - F_0(x - vt, v) \right| dv + \int_{\mathbb{R}^3} F_0(x - vt, v) \left| \int_t^{t_\sigma} g(\tau; t_\sigma, x + \delta, v) d\tau \right| dv \\ &\quad + \int_0^t \int_{\mathbb{R}^3} F_0(x - vt, v) \left| g(\tau; t_\sigma, x + \delta, v) - g(\tau; t, x, v) \right| dv d\tau \\ &\triangleq J_{11} + J_{12} + J_{13}. \end{aligned} \quad (2.5)$$

Let $\tilde{F}(\sigma, \delta) = F_0(x + \delta - vt_\sigma, v) - F_0(x - vt, v)$, it follows from (1.7) that

$$\begin{aligned} J_{11} &\leq \int_{|v| \geq N} \left| \tilde{F}(\sigma, \delta) \right| dv + \int_{|v| \leq N} \left| \tilde{F}(\sigma, \delta) \right| dv \\ &\leq C \|w F_0\|_{L^\infty} \int_{|v| \geq N} (1 + |v|)^{-\beta} dv + \int_{|v| \leq N} \left| \tilde{F}(\sigma, \delta) \right| dv \\ &\leq C(1 + |N|)^{-\beta+3} + \int_{|v| \leq N} \left| \tilde{F}(\sigma, \delta) \right| dv. \end{aligned} \quad (2.6)$$

Fix N large such that $C(1 + |N|)^{-\beta+3} \leq \frac{\varepsilon}{32}$, for $\beta > 3$. Then, from (1.7), there exists $\chi_1 > 0$ depending on (x, t) and ε such that if $\sigma + |\delta| \leq \chi_1$, it holds that

$$J_{11} \leq \frac{\varepsilon}{16}. \quad (2.7)$$

For J_{12} , it follows from (2.2) that

$$\begin{aligned} J_{12} &\leq C \|w F_0\|_{L^\infty} \int_{\mathbb{R}^3} w(v)^{-1} \int_t^{t+\sigma} \int_{\mathbb{R}^3} |v - u|^\gamma w(u)^{-1} \|w F(\tau)\|_{L^\infty} du d\tau dv \\ &\leq C \int_t^{t+\sigma} \|w F(\tau)\|_{L^\infty} d\tau \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} |v - u|^\gamma w(v)^{-1} w(u)^{-1} du dv \\ &\leq C \sigma^{1-\frac{1}{b}} \left(\int_t^{t+\sigma} \|w F(\tau)\|_{L^\infty}^b d\tau \right)^{\frac{1}{b}} \leq C(t) \sigma^{1-\frac{1}{b}} \rightarrow 0 \text{ as } \sigma \rightarrow 0, \end{aligned} \quad (2.8)$$

where we have used the fact that $F \in \mathbf{X}$ and

$$\int_{\mathbb{R}^3} \int_{\mathbb{R}^3} w(v)^{-1} w(u)^{-1} |v - u|^\gamma dudv \leq C < +\infty, \quad \text{for } \beta > 3 + \max\{0, \gamma\}. \quad (2.9)$$

Hence there exists $\chi_2 > 0$ depending on t, ε such that if $\sigma \leq \chi_2$, it holds that

$$J_{12} \leq \frac{\varepsilon}{16}. \quad (2.10)$$

For J_{13} , we note that

$$\begin{aligned} J_{13} &= \int_0^t \int_{\mathbb{R}^3} F_0(x - vt, v) \left| g(\tau; t_\sigma, x + \delta, v) - g(\tau; t_\sigma, x, v) \right| dv d\tau \\ &\quad + \int_0^t \int_{\mathbb{R}^3} F_0(x - vt, v) \left| g(\tau; t_\sigma, x, v) - g(\tau; t, x, v) \right| dv d\tau \\ &\triangleq J_{131} + J_{132}. \end{aligned} \quad (2.11)$$

It holds that

$$\begin{aligned} J_{131} &\leq C \int_0^t \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} w(v)^{-1} w(u)^{-1} |v - u|^\gamma \\ &\quad \times \left| w(u) [F(\tau, x + \delta - v(t_\sigma - \tau), u) - F(\tau, x - v(t_\sigma - \tau), u)] \right| dudvd\tau \\ &= \left\{ \int_{t-\lambda}^t + \int_0^{t-\lambda} \right\} (\cdots) d\tau \triangleq J_{1311} + J_{1312}. \end{aligned} \quad (2.12)$$

From (2.9), one has that

$$J_{1311} \leq C \int_{t-\lambda}^t \|wF(\tau)\|_{L^\infty} d\tau \leq C(t) \lambda^{1-\frac{1}{b}}. \quad (2.13)$$

On the other hand, it holds for some $p > 1$ that

$$\begin{aligned} J_{1312} &\leq C \int_0^{t-\lambda} \left(\int \int \left| w(u) [F(\tau, x + \delta - v(t_\sigma - \tau), u) - F(\tau, x - v(t_\sigma - \tau), u)] \right|^{p_*} dudv \right)^{\frac{1}{p_*}} \\ &\quad \times \left(\int \int w(v)^{-p} w(u)^{-p} |v - u|^{p\gamma} dudv \right)^{\frac{1}{p}} d\tau, \end{aligned} \quad (2.14)$$

where $p_* = \frac{p}{p-1}$. Since $|v - u|^{p\gamma}$ is singular when $-3 < \gamma < 0$, we choose

$$p = 1 + \frac{3 + \gamma}{4(9 - \gamma)}, \quad \text{for } -3 < \gamma \leq 1, \quad (2.15)$$

so that $1 < p \leq \frac{8}{7}$, $p_* = \frac{p}{p-1} \geq 7$, $p\gamma > -3$, $p\frac{3-\gamma}{2} > 3$ and

$$\int_{\mathbb{R}^3} \int_{\mathbb{R}^3} |v - u|^{p\gamma} w(u)^{-p} w(v)^{-p} dudv \leq C < \infty \quad \text{for } \beta > 3 + \max\{0, \gamma\}. \quad (2.16)$$

Changing variable by $y = x - v(t_\sigma - \tau)$, we obtain, for any fixed $\lambda > 0$, that

$$\begin{aligned} J_{1312} &\leq C \lambda^{-\frac{3}{p_*}} \int_0^{t-\lambda} \|wF(\tau)\|_{L^\infty}^{1-\frac{2}{p_*}} \|w(\cdot) [F(\tau, \cdot + \delta, \cdot) - F(\tau, \cdot, \cdot)]\|_{L^2}^{\frac{2}{p_*}} d\tau \\ &\leq C \lambda^{-\frac{3}{p_*}} \left(\int_0^{t-\lambda} \|\cdot\|_{L^\infty}^b d\tau \right)^{\frac{1}{b} - \frac{2}{bp_*}} \left(\int_0^{t-\lambda} \|\cdot\|_{L^2}^{\frac{b}{b-1}} d\tau \right)^{\frac{2(b-1)}{bp_*}} \left(\int_0^{t-\lambda} d\tau \right)^{\tilde{d}} \\ &\leq C(t) \lambda^{-\frac{3}{p_*}} \left(\int_0^{t-\lambda} \|w(\cdot) [F(\tau, \cdot + \delta, \cdot) - F(\tau, \cdot, \cdot)]\|_{L^2}^{\frac{b}{b-1}} d\tau \right)^{\frac{2(b-1)}{bp_*}} \rightarrow 0, \quad \text{as } \delta \rightarrow 0, \end{aligned} \quad (2.17)$$

where $\tilde{d} = 1 - \frac{1}{b} + \frac{4}{bp_*} - \frac{2}{p_*} > 0$ due to $b > 2$, $p_* \geq 7$. From (2.12), (2.13) and (2.17), choosing $\lambda^{1-\frac{1}{b}} = \frac{\varepsilon}{64C(t)}$, then there exists $\chi_3 > 0$ depending only on t, ε such that if $\sigma + |\delta| \leq \chi_3$, one obtains, for $\beta > 3 + \max\{0, \gamma\}$, that

$$J_{131} \leq \frac{\varepsilon}{16}. \quad (2.18)$$

For J_{132} , one notes that

$$\begin{aligned} J_{132} &\leq C \|wF_0\|_{L^\infty} \int_0^t \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} w(v)^{-1} w(u)^{-1} |v - u|^\gamma \\ &\quad \times \left| w(u) [F(\tau, x - v(t_\sigma - \tau), u) - F(\tau, x - v(t - \tau), u)] \right| dudvd\tau \\ &\leq C \left\{ \int_0^t \int_{|v| \geq N} \int_{\mathbb{R}^3} + \int_{t-\lambda}^t \int_{|v| \leq N} \int_{\mathbb{R}^3} + \int_0^{t-\lambda} \int_{|v| \leq N} \int_{\mathbb{R}^3} \right\} (\dots) dudvd\tau \\ &\triangleq J_{1321} + J_{1322} + J_{1323}. \end{aligned} \quad (2.19)$$

It is straightforward to obtain, for $\beta > 3 + \max\{0, \gamma\}$, that

$$J_{1321} \leq C \int_0^t \int_{|v| \geq N} w(v)^{-1} (1 + |v|)^{\max\{0, \gamma\}} dv d\tau \leq C(t) (1 + N)^{-\beta + 3 + \max\{0, \gamma\}} \leq \frac{\varepsilon}{64}, \quad (2.20)$$

and

$$J_{1322} \leq C \int_{t-\lambda}^t \|wF(\tau)\|_{L^\infty} d\tau \leq C(t) \lambda^{1-\frac{1}{b}} \leq \frac{\varepsilon}{64}, \quad (2.21)$$

where N and λ are suitably chosen so that $C(t)(1 + N)^{-\beta + 3 + \max\{0, \gamma\}} \leq \frac{\varepsilon}{64}$ and $\lambda^{1-\frac{1}{b}} = \frac{\varepsilon}{64C(t)}$. Now we estimate J_{1323} . By the definition of $\mathbf{X}$, there exists a smooth compact support function $F^\varepsilon(\cdot, \cdot, \cdot)$ such that

$$\left(\int_0^t \|w(F(\tau, \cdot, \cdot) - F^\varepsilon(\tau, \cdot, \cdot))\|_{L^2}^{\frac{b}{b-1}} d\tau \right)^{\frac{2(b-1)}{bp_*}} \leq \frac{\lambda^{\frac{3}{p_*}} \varepsilon}{C_1}, \quad (2.22)$$

where C_1 will be chosen later. Using (2.22) and the same arguments as in (2.17), for $\beta > 3 + \max\{0, \gamma\}$, we have

$$\begin{aligned} J_{1323} &\leq C \int_0^{t-\lambda} \int_{|v| \leq N} \int_{\mathbb{R}^3} w(v)^{-1} w(u)^{-1} |v - u|^\gamma \\ &\quad \times \left\{ \left| w(u) [F(\tau, x - v(t_\sigma - \tau), u) - F^\varepsilon(\tau, x - v(t_\sigma - \tau), u)] \right| \right. \\ &\quad + \left| w(u) [F(\tau, x - v(t - \tau), u) - F^\varepsilon(\tau, x - v(t - \tau), u)] \right| \\ &\quad \left. + \left| w(u) [F^\varepsilon(\tau, x - v(t_\sigma - \tau), u) - F^\varepsilon(\tau, x - v(t - \tau), u)] \right| \right\} dudvd\tau \\ &\leq C(t) \lambda^{-\frac{3}{p_*}} \left(\int_0^t \|w(F(\tau, \cdot, \cdot) - F^\varepsilon(\tau, \cdot, \cdot))\|_{L^2}^{\frac{b}{b-1}} d\tau \right)^{\frac{2(b-1)}{bp_*}} \\ &\quad + C_{N, \varepsilon} t \sup_{|v| \leq N, (\tau, y, u) \in ((0, T] \times \mathbb{R}^3 \times \mathbb{R}^3)} \left| w(u) [F^\varepsilon(\tau, y - v\sigma, u) - F^\varepsilon(\tau, y, u)] \right| \\ &\leq \frac{C(t)}{C_1} \varepsilon + \frac{\varepsilon}{128} \leq \frac{\varepsilon}{64}, \end{aligned} \quad (2.23)$$

where we have used the fact that

$$\lim_{\sigma \rightarrow 0} C_{N, \varepsilon} t \sup_{|v| \leq N, (\tau, y, u) \in ((0, t] \times \mathbb{R}^3 \times \mathbb{R}^3)} \left| w(u) [F^\varepsilon(\tau, y - v\sigma, u) - F^\varepsilon(\tau, y, u)] \right| = 0, \quad (2.24)$$

and C_1 is chosen large so that (2.23) holds. Thus, from (2.19)-(2.23), there exists $\chi_4 > 0$ depending on t and ε such that if $\sigma \leq \chi_4$, it holds that

$$J_{132} \leq \frac{\varepsilon}{16}. \quad (2.25)$$

Therefore, for $\beta > 3 + \max\{0, \gamma\}$, it follows from (2.5), (2.7), (2.10), (2.11), (2.18) and (2.25) that

$$\left| \int_{\mathbb{R}^3} \left\{ I_1(t + \sigma, x + \delta, v) - I_1(t, x, v) \right\} dv \right| \leq \frac{\varepsilon}{2}, \quad \text{for } \sigma + |\delta| \leq \min\{\chi_1, \chi_2, \chi_3, \chi_4\}. \quad (2.26)$$

Part 2: Estimation on $\int_{\mathbb{R}^3} I_2(t, x, v) dv$.

Note that

$$\begin{aligned} J_2 &\triangleq \left| \int_{\mathbb{R}^3} I_2(t_\sigma, x + \delta, v) - I_2(t, x, v) dv \right| \\ &\leq \left| \int_{\mathbb{R}^3} I_2(t_\sigma, x + \delta, v) - I_2(t_\sigma, x, v) dv \right| + \left| \int_{\mathbb{R}^3} I_2(t_\sigma, x, v) - I_2(t, x, v) dv \right| \\ &\triangleq J_{21} + J_{22}. \end{aligned} \quad (2.27)$$

For J_{21} , it follows from (2.1) that

$$\begin{aligned} J_{21} &\leq \int_0^{t_\sigma} \int_{\mathbb{R}^3} Q_+(F, F)(s, x + \delta - v(t_\sigma - s), v) \int_s^{t_\sigma} |g(\tau; t_\sigma, x + \delta, v) - g(\tau; t_\sigma, x, v)| d\tau dv ds \\ &\quad + \int_0^{t_\sigma} \int_{\mathbb{R}^3} \left| Q_+(F, F)(s, x + \delta - v(t_\sigma - s), v) - Q_+(F, F)(s, x - v(t_\sigma - s), v) \right| dv ds \\ &\triangleq J_{211} + J_{212}. \end{aligned} \quad (2.28)$$

A direct calculation shows that

$$\begin{aligned} Q_+(F, F)(s, y, v) &\leq \int_{\mathbb{R}^3} \int_{\mathbb{S}^2} q(\theta, v - u) w(u')^{-1} w(v')^{-1} |w(u') F(s, y, u')| \cdot |w(v') F(s, y, v')| du d\omega \\ &\leq \|wF(s)\|_{L^\infty}^2 \int_{\mathbb{R}^3} (1 + |v| + |u|)^{-\beta} |v - u|^\gamma du \\ &\leq C(1 + |v|)^{-\beta+3+\max\{0, \gamma\}} \|wF(s)\|_{L^\infty}^2, \quad \text{for } \beta > 3 + \max\{0, \gamma\}, \end{aligned} \quad (2.29)$$

where we have used the fact that

$$w(u')w(v') \geq (1 + |v|^2 + |u|^2)^{\frac{\beta}{2}}, \quad \beta \geq 0, \quad (2.30)$$

and

$$\int_{\mathbb{R}^3} (1 + |v| + |u|)^{-\beta} |v - u|^\gamma du \leq C(1 + |v|)^{-\beta+3+\max\{0, \gamma\}} \quad \text{for } \beta > 3 + \max\{0, \gamma\}. \quad (2.31)$$

Thus, it follows from (2.2) and (2.29) that

$$\begin{aligned} J_{211} &\leq C \int_0^{t_\sigma} \|wF(s)\|_{L^\infty}^2 ds \cdot \int_0^{t_\sigma} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} |v - u|^\gamma w(u)^{-1} \cdot (1 + |v|)^{-\beta+3+\max\{0, \gamma\}} \\ &\quad \times \left| w(u) [F(\tau, x + \delta - v(t_\sigma - \tau), u) - F(\tau, x - v(t_\sigma - \tau), u)] \right| du dv d\tau \\ &\leq C(t) \int_0^{t_\sigma} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} |v - u|^\gamma w(u)^{-1} \cdot (1 + |v|)^{-\beta+3+\max\{0, \gamma\}} \\ &\quad \times \left| w(u) [F(\tau, x + \delta - v(t_\sigma - \tau), u) - F(\tau, x - v(t_\sigma - \tau), u)] \right| du dv d\tau \\ &\leq C(t) \cdot \left\{ \int_{t-\lambda}^{t_\sigma} + \int_0^{t-\lambda} \right\} (\cdots) d\tau \triangleq J_{2111} + J_{2112}. \end{aligned} \quad (2.32)$$

It is straightforward to obtain that

$$\begin{aligned} J_{2111} &\leq C(t) \int_{t-\lambda}^{t_\sigma} \|wF(\tau)\|_{L^\infty} d\tau \int_{\mathbb{R}^3} (1+|v|)^{-\beta+3+2\max\{0,\gamma\}} dv \\ &\leq C(t)(\sigma+\lambda)^{1-\frac{1}{b}}, \quad \text{for } \beta > 6 + 2\max\{0,\gamma\}. \end{aligned} \quad (2.33)$$

On the other hand, for any fixed $\lambda > 0$, it holds that

$$\begin{aligned} J_{2112} &\leq C(t) \int_0^{t-\lambda} \left(\int_{\mathbb{R}^3} \int_{\mathbb{R}^3} |v-u|^{p\gamma} w(u)^{-p} (1+|v|)^{-p\beta+3p+p\max\{0,\gamma\}} dudv \right)^{\frac{1}{p}} \\ &\quad \times \left(\int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \left| w(u) [F(\tau, x+\delta-v(t_\sigma-\tau), u) - F(\tau, x-v(t_\sigma-\tau), u)] \right|^{p_*} dudv \right)^{\frac{1}{p_*}} d\tau \\ &\leq C(t) \lambda^{-\frac{3}{p_*}} \int_0^{t-\lambda} \left(\int_{\mathbb{R}^3} \int_{\mathbb{R}^3} w(u) |F(\tau, y+\delta, u) - F(\tau, y, u)|^{p_*} dudy \right)^{\frac{1}{p_*}} d\tau \\ &\leq C(t) \lambda^{-\frac{3}{p_*}} \left(\int_0^{t-\lambda} \|w(u) [F(\tau, \cdot+\delta, u) - F(\tau, \cdot, u)]\|_{L^2}^{\frac{b}{b-1}} d\tau \right)^{\frac{2(b-1)}{bp_*}} \rightarrow 0 \text{ as } \delta \rightarrow 0, \end{aligned} \quad (2.34)$$

where p is defined in (2.15). Therefore, for $\beta > 6 + 2\max\{0,\gamma\}$, combining (2.32)-(2.34), there exists $\chi_5 > 0$ depending on t and ε such that if $\sigma + |\delta| \leq \chi_5$, it holds that

$$J_{211} \leq \frac{\varepsilon}{32}. \quad (2.35)$$

For J_{212} , we have

$$\begin{aligned} J_{212} &\leq \int_0^{t_\sigma} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \int_{\mathbb{S}^2} q(\theta, v-u) F(s, x+\delta-v(t_\sigma-s), u') \\ &\quad \times \left| F(s, x+\delta-v(t_\sigma-s), v') - F(s, x-v(t_\sigma-s), v') \right| dudvd\omega ds \\ &\quad + \int_0^{t_\sigma} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \int_{\mathbb{S}^2} q(\theta, v-u) F(s, x-v(t_\sigma-s), v') \\ &\quad \times \left| F(s, x+\delta-v(t_\sigma-s), u') - F(s, x-v(t_\sigma-s), u') \right| dudvd\omega ds \\ &\triangleq J_{2121} + J_{2122}. \end{aligned} \quad (2.36)$$

We only consider J_{2121} since J_{2122} can be treated similarly. It follows from (2.30) that

$$\begin{aligned} J_{2121} &\leq \int_0^{t_\sigma} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \int_{\mathbb{S}^2} q(\theta, v-u) w(u')^{-1} w(v')^{-1} \|wF(s)\|_{L^\infty} \\ &\quad \times \left| w(v') [F(s, x+\delta-v(t_\sigma-s), v') - F(s, x-v(t_\sigma-s), v')] \right| dudvd\omega ds \\ &\leq C \int_{t-\lambda}^{t_\sigma} \|wF(s)\|_{L^\infty}^2 \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} |v-u|^\gamma (1+|v|+|u|)^{-\beta} dudv ds \\ &\quad + C \int_0^{t-\lambda} \|wF(s)\|_{L^\infty} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \int_{\mathbb{S}^2} q(\theta, v-u) w(v)^{-\frac{1}{2}} w(u')^{-\frac{1}{2}} w(v')^{-\frac{1}{2}} \\ &\quad \times w(v') \left| F(s, x+\delta-v(t_\sigma-s), v') - F(s, x-v(t_\sigma-s), v') \right| dudvd\omega ds \\ &\triangleq J_{21211} + J_{21212}. \end{aligned} \quad (2.37)$$

It follows from (2.31) that

$$J_{21211} \leq C(t)(\sigma+\lambda)^{1-\frac{2}{b}}. \quad (2.38)$$

For J_{21212} , we follow the idea of [4]. Denote

$$z = u - v, \quad z_{||} = [z \cdot \omega]\omega, \quad z_{\perp} = z - z_{||}, \quad \eta = v + z_{||}, \quad (2.39)$$

one has $u' = v + z_{\perp}$, $v' = v + z_{||}$. A simple calculation yields that

$$q(\theta, z) \leq C|z_{||}| \cdot (|z_{||}| + |z_{\perp}|)^{\gamma-1}. \quad (2.40)$$

Then it follows from (2.30) and (2.40) that

$$\begin{aligned} J_{21212} &\leq C \int_0^{t-\lambda} \|wF(s)\|_{L^\infty} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \int_{\mathbb{S}^2} w(v)^{-\frac{1}{2}} \cdot |z_{||}| \cdot (|z_{||}| + |z_{\perp}|)^{\gamma-1} \cdot w(v + z_{\perp})^{-\frac{1}{2}} w(v + z_{||})^{-\frac{1}{2}} \\ &\quad \times \left| w(v + z_{||}) [F(s, x + \delta - v(t_\sigma - s), v + z_{||}) - F(s, x - v(t_\sigma - s), v + z_{||})] \right| dz_{\perp} d|z_{||}| d\omega dv ds \\ &\quad \text{Noting } |z_{||}|^2 dz_{\perp} d|z_{||}| d\omega = dz_{\perp} d\eta, \\ &\leq C \int_0^{t-\lambda} \|wF(s)\|_{L^\infty} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \int_{z_{\perp}} w(v + z_{\perp})^{-\frac{1}{2}} (|\eta - v| + |z_{\perp}|)^{\gamma-1} w(v)^{-\frac{1}{2}} w(\eta)^{-\frac{1}{2}} |\eta - v|^{-1} \\ &\quad \times \left| w(\eta) [F(s, x + \delta - v(t_\sigma - s), \eta) - F(s, x - v(t_\sigma - s), \eta)] \right| dz_{\perp} d\eta dv ds. \end{aligned} \quad (2.41)$$

We note, for $\beta > 4$, that

$$\begin{aligned} &\int_{z_{\perp}} w(v + z_{\perp})^{-\frac{1}{2}} (|\eta - v| + |z_{\perp}|)^{\gamma-1} dz_{\perp} \leq |\eta - v|^{\frac{\gamma-1}{2}} \int_{z_{\perp}} w(v + z_{\perp})^{-\frac{1}{2}} |z_{\perp}|^{\frac{\gamma-1}{2}} dz_{\perp} \\ &\leq |\eta - v|^{\frac{\gamma-1}{2}} \left\{ \int_{|z_{\perp}| \leq 1} |z_{\perp}|^{\frac{\gamma-1}{2}} dz_{\perp} + \int_{|z_{\perp}| \geq 1} (1 + |v + z_{\perp}|)^{-\frac{\beta}{2}} dz_{\perp} \right\} \leq C |\eta - v|^{\frac{\gamma-1}{2}}, \end{aligned} \quad (2.42)$$

which, together with (2.15), yields, for $\beta > 6$ and any fixed $\lambda > 0$, that

$$\begin{aligned} J_{21212} &\leq C \int_0^{t-\lambda} \|wF(s)\|_{L^\infty} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} w(v)^{-\frac{1}{2}} w(\eta)^{-\frac{1}{2}} |\eta - v|^{\frac{\gamma-3}{2}} \\ &\quad \times \left| w(\eta) [F(s, x + \delta - v(t_\sigma - s), \eta) - F(s, x - v(t_\sigma - s), \eta)] \right| d\eta dv ds \\ &\leq C \lambda^{-\frac{3}{p^*}} \int_0^t \|wF(s)\|_{L^\infty}^{2-\frac{2}{p^*}} \|w(F(s, \cdot + \delta, \cdot) - F(s, \cdot, \cdot))\|_{L^2}^{\frac{2}{p^*}} \\ &\quad \times \left(\int_{\mathbb{R}^3} \int_{\mathbb{R}^3} w(v)^{-\frac{p}{2}} w(\eta)^{-\frac{p}{2}} |\eta - v|^{p\frac{\gamma-3}{2}} d\eta dv \right)^{\frac{1}{q}} ds \\ &\leq C(t) \lambda^{-\frac{3}{p^*}} \left(\int_0^t \|w(F(s, \cdot + \delta, \cdot) - F(s, \cdot, \cdot))\|_{L^2}^{\frac{b}{b-1}} ds \right)^{\frac{2(b-1)}{bp^*}} \rightarrow 0, \text{ as } \delta \rightarrow 0. \end{aligned} \quad (2.43)$$

Combining (2.36)-(2.38) and (2.43), for $\beta > 6 + \max\{0, \gamma\}$, there exists $\chi_6 > 0$ depending on t and ε such that if $\sigma + |\delta| \leq \chi_6$, one has that

$$J_{212} \leq \frac{\varepsilon}{32}. \quad (2.44)$$

It remains to estimate J_{22} . A direct calculation shows that

$$\begin{aligned} J_{22} &\leq \int_t^{t+\sigma} \int_{\mathbb{R}^3} Q_+(F, F)(s, x - v(t_\sigma - s), v) dv ds \\ &\quad + \int_0^t \int_{\mathbb{R}^3} Q_+(F, F)(s, x - v(t - s), v) \int_t^{t+\sigma} g(\tau; t_\sigma, x, v) d\tau dv ds \\ &\quad + \int_0^t \int_{\mathbb{R}^3} Q_+(F, F)(s, x - v(t - s), v) \int_s^t \left| g(\tau; t_\sigma, x, v) - g(\tau; t, x, v) \right| d\tau dv ds \\ &\quad + \int_0^t \int_{\mathbb{R}^3} \left| Q_+(F, F)(s, x - v(t_\sigma - s), v) - Q_+(F, F)(s, x - v(t - s), v) \right| dv ds \\ &\triangleq J_{221} + J_{222} + J_{232} + J_{224}. \end{aligned} \quad (2.45)$$

It follows from (2.29), for $\beta > 6 + \max\{0, \gamma\}$, that

$$J_{221} \leq C \int_t^{t+\sigma} \|wF(s)\|_{L^\infty}^2 ds \leq C(t) \sigma^{1-\frac{2}{b}} \rightarrow 0, \quad \text{as } \sigma \rightarrow 0, \quad (2.46)$$

and

$$\begin{aligned} J_{222} &\leq \int_0^t \int (1+|v|)^{-\beta+3+\max\{0,\gamma\}} \|wF(s)\|_{L^\infty}^2 \int_t^{t+\sigma} \int |v-u|^\gamma w(u)^{-1} \|wF(\tau)\|_{L^\infty} dud\tau dv ds \\ &\leq C(t) \int_t^{t+\sigma} \|wF(s)\|_{L^\infty} ds \leq C(t) \sigma^{1-\frac{1}{b}} \rightarrow 0, \quad \text{as } \sigma \rightarrow 0. \end{aligned} \quad (2.47)$$

Using the same arguments as in (2.20)-(2.23), we have that if σ is small,

$$\begin{aligned} J_{223} &\leq C \int_0^t \|wF(s)\|_{L^\infty}^2 ds \int_0^t \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} |v-u|^\gamma w(u)^{-1} (1+|v|)^{-\beta+3+\max\{0,\gamma\}} \\ &\quad \times \left| w(u) [F(\tau, x-v(t+\sigma-\tau), u) - F(\tau, x-v(t-\tau), u)] \right| dudvd\tau \\ &\leq C(t) \int_{t-\lambda}^t \|wF(\tau)\|_{L^\infty} d\tau + C(t) \int_{|v|\geq N} \int_{\mathbb{R}^3} |v-u|^\gamma w(u)^{-1} (1+|v|)^{-\beta+3+\max\{0,\gamma\}} dudv \\ &\quad + C(t) \int_0^{t-\lambda} \int_{|v|\leq N} \int_{\mathbb{R}_u^3} |v-u|^\gamma w(u)^{-1} (1+|v|)^{-\beta+3+\max\{0,\gamma\}} \\ &\quad \times w(u) \left| F(\tau, x-v(t+\sigma-\tau), u) - F(\tau, x-v(t-\tau), u) \right| dudvd\tau \leq \frac{\varepsilon}{16}. \end{aligned} \quad (2.48)$$

For J_{224} , one notes that

$$\begin{aligned} J_{224} &\leq \int_0^t \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \int_{\mathbb{S}^2} q(\theta, v-u) F(s, x-v(t+\sigma-s), u') \\ &\quad \times |F(s, x-v(t+\sigma-s), v') - F(s, x-v(t-s), v')| dudvd\omega ds \\ &\quad + \int_0^t \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \int_{\mathbb{S}^2} q(\theta, v-u) F(s, x-v(t-s), v') \\ &\quad \times |F(s, x-v(t+\sigma-s), u') - F(s, x-v(t-s), u')| dudvd\omega ds \\ &\triangleq J_{2241} + J_{2242}. \end{aligned} \quad (2.49)$$

We only consider J_{2241} since the argument for J_{2242} is similar. It holds that

$$\begin{aligned} J_{2241} &\leq C \int_{t-\lambda}^t \|wF(s)\|_{L^\infty}^2 ds \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} |v-u|^\gamma (1+|v|+|u|)^{-\beta} dudvd s \\ &\quad + \int_0^{t-\lambda} \|wF(s)\|_{L^\infty}^2 ds \int_{|v|\geq N} \int_{\mathbb{R}^3} |v-u|^\gamma (1+|v|+|u|)^{-\beta} dudv \\ &\quad + \int_0^{t-\lambda} \|wF(s)\|_{L^\infty} \int_{|v|\leq N} \int_{\mathbb{R}^3} \int_{\mathbb{S}^2} q(\theta, v-u) w(u')^{-1} w(v')^{-1} \\ &\quad \times \left| w(v') [F(s, x-v(t+\sigma-s), v') - F(s, x-v(t-s), v')] \right| dudvd\omega ds \\ &\triangleq J_{22411} + J_{22412} + J_{22413}. \end{aligned} \quad (2.50)$$

By the same arguments as in (2.20) and (2.21), one can get that

$$J_{22411} + J_{22412} \leq C(t) \lambda^{1-\frac{2}{b}} + C(t) (1+N)^{-\beta+6+\max\{0,\gamma\}} \leq \frac{\varepsilon}{16}. \quad (2.51)$$

For J_{22413} , we notice that

$$\begin{aligned}
J_{22413} &\leq \int_0^{t-\lambda} \|wF(s)\|_{L^\infty} \int_{|v|\leq N} \int_{\mathbb{R}^3} \int_{\mathbb{S}^2} q(\theta, v-u) w(v)^{-\frac{1}{2}} w(u')^{-\frac{1}{2}} w(v')^{-\frac{1}{2}} \\
&\quad \times \left\{ \left| w(v') [F(s, x-v(t+\sigma-s), v') - F^\varepsilon(s, x-v(t+\sigma-s), v')] \right| \right. \\
&\quad \left. + \left| w(v') [F(s, x-v(t-s), v') - F^\varepsilon(s, x-v(t-s), v')] \right| \right\} dudvd\omega ds \\
&\quad + \int_0^{t-\lambda} \|wF(s)\|_{L^\infty} \int_{|v|\leq N} \int_{\mathbb{R}^3} \int_{\mathbb{S}^2} q(\theta, v-u) w(u')^{-1} w(v')^{-1} \\
&\quad \times \left| w(v') [F^\varepsilon(s, x-v(t+\sigma-s), v') - F^\varepsilon(s, x-v(t-s), v')] \right| dudvd\omega ds \\
&\triangleq J_{224131} + J_{224132}.
\end{aligned} \tag{2.52}$$

It follows from (2.22) and the same arguments as in (2.41)-(2.43), for $\beta > 6$, that

$$\begin{aligned}
J_{224131} &\leq \lambda^{-\frac{3}{p_*}} \int_0^t \|wF(s)\|_{L^\infty}^{\frac{2-\frac{2}{p_*}}{p_*}} \|w[F(\tau, \cdot, \cdot) - F^\varepsilon(\tau, \cdot, \cdot)]\|_{L^2}^{\frac{2}{p_*}} d\tau \\
&\leq C(t) \lambda^{-\frac{3}{p_*}} \left(\int_0^t \|w[F(\tau, \cdot, \cdot) - F^\varepsilon(\tau, \cdot, \cdot)]\|_{L^2}^{\frac{b}{b-1}} d\tau \right)^{\frac{2(b-1)}{bp_*}} \leq \frac{\varepsilon}{32}.
\end{aligned} \tag{2.53}$$

For J_{224132} , one can obtain, for $\beta > 3 + \max\{0, \gamma\}$, that

$$J_{224132} \leq C_{N,\varepsilon} t \sup_{|v|\leq N, (\tau, y, u) \in ((0, T] \times \mathbb{R}^3 \times \mathbb{R}^3)} w(u) \left| F^\varepsilon(\tau, y - v\sigma, u) - F^\varepsilon(\tau, y, u) \right| \rightarrow 0, \text{ as } \sigma \rightarrow 0. \tag{2.54}$$

Combining (2.45)-(2.54), for $\beta > 6 + 2\max\{0, \gamma\}$, there exists $\chi_7 > 0$ depending on t and ε such that if $\sigma \leq \chi_7$, one has that

$$J_{22} \leq \frac{\varepsilon}{32}. \tag{2.55}$$

Thus it follows from (2.27), (2.28), (2.35), (2.44) and (2.55), $\beta > 6 + 2\max\{0, \gamma\}$, that

$$\left| \int_{\mathbb{R}^3} \left\{ I_2(t+\sigma, x+\delta, v) - I_2(t, x, v) \right\} dv \right| \leq \frac{\varepsilon}{2}, \text{ for } \sigma + |\delta| \leq \min\{\chi_5, \chi_6, \chi_7\}. \tag{2.56}$$

Taking $\chi \triangleq \min\{\chi_1, \chi_2, \chi_3, \chi_4, \chi_5, \chi_6, \chi_7\}$, combining (2.1), (2.26) and (2.56), for $\beta > 6 + 2\max\{0, \gamma\}$, we have

$$\left| \int_{\mathbb{R}^3} \left\{ F(t+\sigma, x+\delta, v) - F(t, x, v) \right\} dv \right| \leq \varepsilon, \text{ for } \sigma + |\delta| \leq \chi. \tag{2.57}$$

That is $\int F(t, x, v) dv$ is continuous function of $(x, t) \in \mathbb{R}^3 \times (0, \infty)$. In the same way, for $\beta > 8 + 2\max\{0, \gamma\}$, one can prove $\int vF(t, x, v)$ and $\int |v|^2 F(t, x, v) dv$ are also continuous functions of $(x, t) \in \mathbb{R}^3 \times (0, \infty)$. Thus the proof of Theorem 1.1 is completed. $\square$

Proof of Theorem 1.2. For any fixed $0 < t_1 < T + \infty$, let $(x, t) \in \mathbb{R} \times [t_1, T]$. The uniform continuity of density in $(x, t) \in \mathbb{R}^3 \times [t_1, T]$ is equivalent to prove that for any small $\varepsilon > 0$, there exists $\chi > 0$ depending on T, t_1^{-1} and ε such that if $|\sigma| + |\delta| \leq \chi$, it holds that

$$\left| \int_{\mathbb{R}^3} F(t+\sigma, x+\delta, v) - F(t, x, v) dv \right| \leq \varepsilon. \tag{2.58}$$

Without loss of generality, we assume $\sigma \geq 0$. As in (2.5), one obtains that

$$\begin{aligned}
K_1 &\triangleq \left| \int_{\mathbb{R}^3} (I_1(t_\sigma, x + \delta, v) - I_1(t, x, v)) dv \right| \\
&\leq \int_{\mathbb{R}^3} \left| F_0(x + \delta - vt_\sigma, v) - F_0(x - vt, v) \right| dv + \int_{\mathbb{R}^3} F_0(x - vt, v) \left| \int_t^{t_\sigma} g(\tau, t_\sigma, x + \delta, v) d\tau \right| dv \\
&\quad + \int_0^t \int_{\mathbb{R}^3} F_0(x - vt, v) \left| g(\tau; t_\sigma, x + \delta, v) - g(\tau; t, x, v) \right| dv d\tau \\
&\triangleq K_{11} + K_{12} + K_{13}.
\end{aligned} \tag{2.59}$$

By the same arguments as in (2.6), one has, for $\beta > 3$, that

$$K_{11} \leq \frac{\varepsilon}{32} + \sup_{(t, x) \in [t_1, T] \times \mathbb{R}^3} \int_{|v| \leq N} \left| F_0(x + \delta - v(t + \sigma), v) - F_0(x - vt, v) \right| dv. \tag{2.60}$$

It follows from (1.15) that there exists $\chi_1 > 0$ depending on t_1^{-1}, T and ε such that if $\sigma + |\delta| \leq \chi_1$, one has that

$$\sup_{(t, x) \in [t_1, T] \times \mathbb{R}^3} \int_{|v| \leq N} \left| F_0(x + \delta - v(t + \sigma), v) - F_0(x - vt, v) \right| dv \leq \frac{\varepsilon}{32}, \tag{2.61}$$

which, together with (2.60), for $\beta > 3$, that

$$K_{11} \leq \frac{\varepsilon}{16}, \text{ for } \sigma + |\delta| \leq \chi_1. \tag{2.62}$$

On the other hand, by the same arguments as in (2.8)-(2.25), one can prove that there exists $\chi_2 > 0$ depending on t_1^{-1}, T and ε such that if $\sigma + |\delta| \leq \chi_2$, it holds, for $\beta > 6 + 2 \max\{0, \gamma\}$, that

$$K_{12} + K_{13} \leq \frac{\varepsilon}{16}. \tag{2.63}$$

As in (2.27)-(2.56), one can prove that there exists $\chi_3 > 0$ depending on t_1^{-1}, T and ε such that if $\sigma + |\delta| \leq \chi_3$, one obtains, for $\beta > 6 + 2 \max\{0, \gamma\}$, that

$$K_2 \triangleq \left| \int_{\mathbb{R}^3} I_2(t + \sigma, x + \delta, v) - I_2(t, x, v) dv \right| \leq \frac{\varepsilon}{16}. \tag{2.64}$$

Taking $\chi = \min\{\chi_1, \chi_2, \chi_3\}$, then we have, for $\beta > 6 + 2 \max\{0, \gamma\}$, that

$$\left| \int_{\mathbb{R}^3} F(t + \sigma, x + \delta, v) - F(t, x, v) dv \right| \leq \varepsilon \text{ for } \sigma + |\delta| \leq \chi. \tag{2.65}$$

Thus $\int_{\mathbb{R}^3} F(t, x, v) dv$ is uniformly continuous in $(x, t) \in \mathbb{R}^3 \times [t_1, T]$. In the same way, for $\beta > 8 + 2 \max\{0, \gamma\}$, one can prove $\int v F(t, x, v) dv$ and $\int |v|^2 F(t, x, v) dv$ are also uniformly continuous in $(x, t) \in \mathbb{R}^3 \times [t_1, T]$. Therefore the proof of Theorem 1.2 is completed. $\square$

3 Appendix

We shall prove that the examples given in (1.8) and (1.11) satisfy the condition (1.7) and (1.15). Here we only give the proof for (1.8) since the case for (1.11) is similar.

Using (1.9), it is obvious that $\|wF_0\|_{L^\infty} \leq C < \infty$. Next we prove the second part of (1.7). We only need to prove that for any small $\varepsilon > 0$, there exists $\chi > 0$ depending on t, ε such that if $\sigma + |\delta| \leq \chi$, it holds that

$$\int_{\mathbb{R}^3} |F_0(x + \delta - v(t + \sigma), v) - F_0(x - vt, v)| dv \leq \varepsilon. \quad (3.1)$$

We note that, for any fixed $t > 0$, that

$$\begin{aligned} & \int_{\mathbb{R}^3} |F_0(x + \delta - v(t + \sigma), v) - F_0(x - vt, v)| dv \\ & \leq \int_{\mathbb{R}^3} |F_0(x + \delta - v(t + \sigma), v) - F_0(x - v(t + \sigma), v)| dv \\ & \quad + \int_{\mathbb{R}^3} |F_0(x - v(t + \sigma), v) - F_0(x - vt, v)| dv \\ & \triangleq L_1 + L_2. \end{aligned} \quad (3.2)$$

Without loss of generality, we assume $\sigma \geq 0$. By changing $y = x - vt_\sigma$ and (1.10), we have

$$\begin{aligned} L_1 & \leq C \left\{ \int_{\mathbb{R}^3} |(\rho_0(x + \delta - vt_\sigma) - \rho_0(x - vt_\sigma))^p + (u_0(x + \delta - vt_\sigma) - u_0(x - vt_\sigma))^p \right. \\ & \quad \left. + (\theta_0(x + \delta - vt_\sigma) - \theta_0(x - vt_\sigma))^p dv \right\}^{\frac{1}{p}} \\ & \leq C t_\sigma^{-\frac{3}{p}} \left\{ \|\rho_0(\cdot + \delta) - \rho_0(\cdot)\|_{L^p} + \|u_0(\cdot + \delta) - u_0(\cdot)\|_{L^p} + \|\theta_0(\cdot + \delta) - \theta_0(\cdot)\|_{L^p} \right\} \\ & \rightarrow 0, \quad \text{as } \sigma, \delta \rightarrow 0, \end{aligned} \quad (3.3)$$

where we have used the continuity of translations on L^p . Thus there exists $\chi_1 > 0$ depending on $t > 0$ and $\varepsilon > 0$ such that if $\sigma + |\delta| \leq \chi_1$, it holds that

$$L_1 \leq \frac{\varepsilon}{2}. \quad (3.4)$$

For L_2 , it is straightforward to obtain that

$$\begin{aligned} L_2 & \leq C \int_{\mathbb{R}^3} \left(|(\rho_0(x - vt - v\sigma) - \rho_0(x - vt))| + |u_0(x - vt - v\sigma) - u_0(x - vt)| \right. \\ & \quad \left. + |\theta_0(x - vt - v\sigma) - \theta_0(x - vt)| \right) e^{-\frac{|v|^2}{8C}} dv. \end{aligned} \quad (3.5)$$

Noting that $\rho_0 - 1 \in L^p$ and $0 \leq \rho_0(x) \leq \hat{C}$, there exists a smooth compact support function ρ_0^ε such that

$$\|\rho_0 - \rho_0^\varepsilon\|_{L^p} \leq \frac{\varepsilon}{C_2(t)}, \quad (3.6)$$

where $C_2(t)$ is a large constant. Hence one obtains that

$$\begin{aligned}
& \int_{\mathbb{R}^3} |\rho_0(x - vt - v\sigma) - \rho_0(x - vt)| e^{-\frac{|v|^2}{8C}} dv \\
& \leq \int_{|v| \geq N} |\rho_0(x - vt - v\sigma) - \rho_0(x - vt)| e^{-\frac{|v|^2}{8C}} dv + \int_{|v| \leq N} |\rho_0(x - vt - v\sigma) - \rho_0(x - vt)| e^{-\frac{|v|^2}{8C}} dv \\
& \leq C e^{-\frac{N^2}{16C}} + \int_{|v| \leq N} |\rho_0(x - vt - v\sigma) - \rho_0^\varepsilon(x - vt - v\sigma)| e^{-\frac{|v|^2}{8C}} dv \\
& \quad + \int_{|v| \leq N} |\rho_0(x - vt) - \rho_0^\varepsilon(x - vt)| e^{-\frac{|v|^2}{8C}} dv + \int_{|v| \leq N} |\rho_0^\varepsilon(x - vt - v\sigma) - \rho_0^\varepsilon(x - vt)| e^{-\frac{|v|^2}{8C}} dv \\
& \leq C e^{-\frac{N^2}{16C}} + C t^{-\frac{3}{p}} \|\rho_0 - \rho_0^\varepsilon\|_{L^p} + \int_{|v| \leq N} |\rho_0^\varepsilon(x - vt - v\sigma) - \rho_0^\varepsilon(x - vt)| e^{-\frac{|v|^2}{8C}} dv \\
& \leq \frac{C t^{-\frac{3}{p}}}{C_2(t)} \varepsilon + C e^{-\frac{N^2}{16C}} + C \sup_{y \in \mathbb{R}^3, |v| \leq N} \|\rho_0^\varepsilon(y + v\sigma) - \rho_0^\varepsilon(y)\|_{L^\infty} \leq \frac{\varepsilon}{16}.
\end{aligned} \tag{3.7}$$

Similarly, we can deal with the remaining terms of (3.5). Therefore, there exists $\chi_2 > 0$ depending on t and ε such that if $\sigma \leq \chi_2$, it holds that

$$L_2 \leq \frac{\varepsilon}{4}. \tag{3.8}$$

Combining (3.4) and (3.8), there exists $\chi = \min\{\chi_1, \chi_2\}$ depending on t and ε such that (3.1) holds if $\sigma + |\delta| \leq \chi$. Therefore the condition (1.7) holds for the example given in (1.8).

Finally, for any fixed $0 < t_1 < T < \infty$, from (3.4) and (3.8), it is not difficult to show that there exists $\chi > 0$ depending on t_1^{-1}, T and ε such that if $\sigma + |\delta| \leq \chi$, it holds that

$$\sup_{(x,t) \in \mathbb{R}^3 \times [t_1, T]} \int_{\mathbb{R}^3} |F_0(x + \delta - v(t + \sigma), v) - F_0(x - vt, v)| dv \leq \varepsilon. \tag{3.9}$$

Thus the example given in (1.8) also satisfies the condition (1.15).

Acknowledgments. Feimin Huang is partially supported by by National Center for Mathematics and Interdisciplinary Sciences, AMSS, CAS and NSFC Grant No.11371349. Yong Wang is partially supported by NSFC Grant No. 11401565.

References

- [1] R. Alexandre, Y. Morimoto, S. Ukai, C.-J. Xu, T. Yang, *Smoothing effect of weak solutions for the spatially homogeneous Boltzmann equation without angular cutoff*. Kyoto J. Math. **52**(2012), no. 3, 433-463.
- [2] R. Alexandre, Y. Morimoto, S. Ukai, C.-J. Xu, T. Yang, *Regularizing effect and local existence for the non-cutoff Boltzmann equation*. Arch. Ration. Mech. Anal. **198** (2010), no. 1, 39-123.
- [3] K. Aoki, C. Bardos, C. Dogbe, F. Golse, *A note on the propagation of boundary induced discontinuities in kinetic theory*. Math. Models Methods Appl. Sci. **11**(9)(2001), 1581-1595.
- [4] N. Bellomo, A. Palczewski, G. Toscani, *Mathematical topics in nonlinear kinetic theory*, World Scientific Publishing Co., Singapore, 1988.

- [5] N. Bellomo, G. Toscani, *On the Cauchy problem for the nonlinear Boltzmann equation global existence uniqueness and asymptotic stability*, J. Math. Phys, **26**(1984), 334-338.
- [6] L. Bernis, L. Desvillettes, *Propagation of singularities for classical solutions of the Vlasov-Poisson- Boltzmann equation*. Discrete Contin. Dyn. Syst. **24**(2009), 13-33.
- [7] L. Boudin and L. Desvillettes, *On the singularities of the global small solution soft the full Boltzmann equation*, Monatsch. Math. **131**(2000), 91-108.
- [8] M. Cannone, G. Karch, *Infinite energy solutions to the homogeneous Boltzmann equation*, Comm. Pure Appl. Math. **63**(2010), 747-778.
- [9] E.A. Carlen, E. Gabetta, G. Toscani, *Propagation of smoothness and the rate of exponential convergence to equilibrium for a spatially homogeneous Maxwellian gas*, Comm. Math. Phys. 199 (1999) 521-546.
- [10] C. Cercignani, R. Illner, M. Pulvirenti, *The Mathematical Theory of Dilute Gases*, Springer-Verlag, New York, 1994.
- [11] R. J. Duan, *On the Cauchy problem for the Boltzmann equation in the whole space: Global existence and uniform stability in $L^2_\xi(H^N_x)$* , J. Differential Equations. **244**(2008), 3204-3234
- [12] R.J. DiPerna, P.-L. Lions, *On the Cauchy problem for Boltzmann equation: Global existence and weak stability*, Ann. of Math. **130** (1989), 321-366.
- [13] L. Desvillettes, B. Wennberg, *Smoothness of the solution of the spatially homogeneous Boltzmann equation without cutoff*, Comm. Partial Diff. Eqns. **29** (2004), 133-155.
- [14] R.J. Duan, M.R. Li, T. Yang, *Propagation of singularities in the solutions to the Boltzmann equation near equilibrium*, Math. Models Methods Appl. Sci. **18**(2008), 1093-1114.
- [15] R.T. Glassey, *The cauchy problem in kinetic theory*, Society for Industrial and Applied Mathematics (SIAM), Philadelphia, 1996.
- [16] F. Golse, P.L. Lions, B. Perthame, R. Sentis, *Regularity of the moments of the solution of a transport equation*, J. Funct. Anal. 76 (1988), no. 1, 110125.
- [17] Y. Guo, *The Boltzmann equation in the whole space*, Indiana Univ. Math. J. **53**(2004), 1081-1094.
- [18] Y. Guo, *Bounded solutions for the Boltzmann equation*, Quart. Appl. Math. **68** (2010), no.1, 143-148.
- [19] K. Hamdache, *Existence in the large and asymptotic behaviour for the Boltzmann equation*, Jpn. J. Appl. Math. **2**(1985), 1-15.
- [20] D. Hoff and T.P. Liu, *The inviscid limit for the Navier-Stokes equations of compressible isentropic flow with shock data*, Indiana Univ. Math. J. **38**(1989), no.(4), 861-915.
- [21] R. Illner, M. Shinbrot, *Global existence for a rare gas in an infinite vacuum*, Comm. Math. Phys. **95** (1984), 117-126.
- [22] K. Imai, T.Nishida, *Global solutions to the initial value problem for the nonlinear Boltzmann equation*, Publ. RIMS Kyoto Univ. **12** (1976), 229-239.
- [23] S. Kaniel, M. Shinbrot, *The Boltzmann equation I: Uniqueness and global existence*, Comm. Math. Phys. **58** (1978) 65-84.

- [24] C. Kim, *Formation and propagation of discontinuity for Boltzmann equation in non-convex domain* Comm. Math. Phys. **308** (2011) 641-701.
- [25] S. Mischler, B. Perthame, *Boltzmann equation with infinite energy: renormalized solutions and distributional solutions for small initial data and initial data close to a Maxwellian*, SIAM. J. Math Anal **28**(1997), 1015-1027.
- [26] Y. Morimoto, S.K. Wang, T. Yang, *A new characterization and global regularity of infinite energy solutions to the homogeneous Boltzmann equation*, J. Math. Pures Appl. **103**(2015), 809-829.
- [27] C. Mouhot, C. Villani, *Regularity theory for the spatially homogeneous Boltzmann equation with cut-off*, Arch. Rational. Mech. Anal. **173**(2004), 169-212.
- [28] J. Polewczak, *Classical solutions of the Boltzmann equation in all 3: asymptotic behavior of solutions*, J. Stat. Phys, **50** (1988), 61-632.
- [29] S. Ukai and T. Yang, *The Boltzmann equation in the space $L^2 \cap L^\infty_\beta$: Global and time-periodic solutions*, Anal. Appl. **4** (2006), 263-310.
- [30] C. Villani, *A review of mathematical topics in collisional kinetic theory. Handbook of mathematical fluid dynamics*, Vol. I, 71-305. North-Holland, Amsterdam, 2002.